



King's Research Portal

DOI:

[10.1016/j.ejogrb.2017.11.012](https://doi.org/10.1016/j.ejogrb.2017.11.012)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Moore, A. P., D'Amico, M. I., Cooper, N. A. M., & Thangaratinam, S. (2018). Designing a lifestyle intervention to reduce risk of type 2 diabetes in postpartum mothers following gestational diabetes: An online survey with mothers and health professionals. *European Journal of Obstetrics Gynecology and Reproductive Biology*, 220, 106-112. <https://doi.org/10.1016/j.ejogrb.2017.11.012>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Title: Designing a lifestyle intervention to reduce risk of type 2 diabetes in postpartum mothers following gestational diabetes: An online survey with mothers and health professionals

Authors: Amanda P. Moore, Maria I. D'Amico, Natalie A.M. Cooper, Shakila Thangaratinam



PII: S0301-2115(17)30522-5
DOI: <https://doi.org/10.1016/j.ejogrb.2017.11.012>
Reference: EURO 10117

To appear in: *EURO*

Received date: 14-7-2017
Revised date: 17-10-2017
Accepted date: 13-11-2017

Please cite this article as: Moore Amanda P, D'Amico Maria I, Cooper Natalie AM, Thangaratinam Shakila. Designing a lifestyle intervention to reduce risk of type 2 diabetes in postpartum mothers following gestational diabetes: An online survey with mothers and health professionals. *European Journal of Obstetrics and Gynecology and Reproductive Biology* <https://doi.org/10.1016/j.ejogrb.2017.11.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Designing a lifestyle intervention to reduce risk of type 2 diabetes in postpartum mothers following gestational diabetes: An online survey with mothers and health professionals

Running header: **lifestyle intervention to reduce risk of type 2 diabetes in postpartum mothers following gestational diabetes**

Amanda P. Moore, MSc, BSc (Hons), Maria I. D'Amico, BSc (Hons), Dr Natalie A.M. Cooper, PhD, MBChB, Professor Shakila Thangaratinam, PhD, MRCOG

From the Diabetes and Nutritional Sciences Division, Faculty of Life Sciences & Medicine, King's College London, UK (Amanda Moore); Women's Health Research Unit, Barts and the London School of Medicine and Dentistry, Queen Mary University London, London, UK (Maria D'Amico, Dr Natalie Cooper, Professor Shakila Thangaratinam), Multidisciplinary Evidence Synthesis Hub (mEsh), Barts and the London School of Medicine and Dentistry, Queen Mary University London, London, UK (Professor Shakila Thangaratinam).

Corresponding author:

Maria I. D'Amico, Women's Health Research Unit, Centre for Primary Care and Public Health, Blizard Institute, Barts and The London School of Medicine and Dentistry, London, E1 2AB
Tel: +44 20 7882 5881, Email: m.damico@qmul.ac.uk

Abstract

Objective

The aim of this study was to identify what components of a postpartum lifestyle intervention would engage postpartum mothers who had a diagnosis of gestational diabetes.

Study design

Two online surveys were conducted, one involving postnatal mothers with GDM (n=83), and a second for health professionals (n=46).

Results

Seventy-eight percent of mothers were aware that healthy eating, exercise and weight management were all important to reduce risk of subsequent type 2 diabetes. However, 80% of women in this survey were not ready to engage in a postpartum lifestyle intervention within the first 6 months of giving birth; in contrast 52% of health professionals recommended they should be engaged in the first six weeks. Group sessions were the most commonly chosen format to deliver an intervention (30%). A community setting was preferred to a medical one. Mothers wanted recipe ideas (95%) in preference to general dietary advice (76%) or cooking skills courses (39%). Walking was the main form of exercise for 79% of mothers in this sample. Women highlighted difficulty in focusing on their own health goals because of competing demands of looking after a baby (41% agreed, Median 3, IQR 2), tiredness (65% agreed, Md 4, IQR 1) and the need for childcare (64% agreed, Md 4, IQR 2).

Conclusion

A walking programme, recipe ideas and weight monitoring may be useful components when designing a postpartum lifestyle intervention. Barriers to engagement are evident and the intervention should allow women to engage at a time that is appropriate for them.

Keywords: Gestational diabetes; Weight; Postpartum; survey; Pregnancy; Women's Health; Type 2 diabetes, lifestyle intervention

Introduction

The rise in obesity has led to an increase in the number of women diagnosed with gestational diabetes (GDM), a condition characterised by glucose intolerance first detected during pregnancy¹. Approximately 40% of women with GDM progress to type 2 diabetes mellitus in the first five years following delivery². Furthermore, women with GDM are at risk of cardiovascular disease, ophthalmic morbidity, female malignancies and high rates of obesity in their offspring³⁻⁵. A diagnosis of GDM presents an opportunity to identify women at increased risk of developing type 2 diabetes, and to offer them interventions that could reduce this risk.

Lifestyle interventions involving weight-loss, diet and exercise are effective in reducing the progression to type 2 diabetes mellitus in non-pregnant, pre-diabetes populations⁶.

Randomised trials on effects of intervention after a diagnosis of GDM have shown varied effects in reducing the risk factors for diabetes, alongside breastfeeding and pharmacological intervention, lifestyle based interventions, show potential to reduce progression to type 2 diabetes mellitus⁷, however robust trial data for the effectiveness of lifestyle intervention is limited to date and comes primarily from Asia and the US⁸.

In the postpartum period, competing needs of a young baby can limit motivation for weight loss⁹, healthy eating¹⁰ and intensive lifestyle intervention^{11, 12} and can make it difficult to

engage women in interventions. It has also been reported that women who had GDM may not be aware, or concerned about their diabetes risk^{13, 14} which may influence their postpartum health behaviours.

This study has been undertaken to inform the design of a lifestyle intervention for postpartum GDM mothers in the UK. The objectives were to understand what would be the best timing to recruit to an intervention, what mothers and health professionals thought the programme should include and in what format it should be delivered. The authors also aimed to understand the influence of barriers to engagement to positive postpartum health behaviours in this population and whether health professionals who advised women with GDM about weight and post-natal health, would support referring women into a lifestyle intervention.

Methods

Design and setting: Two on-line surveys were carried out between February 2015 and January 2016, one involving postnatal women in UK with a diagnosis of GDM in a recent pregnancy, and a second involving healthcare professionals caring for women with GDM. The surveys were hosted on-line using an on-line survey tool (Survey Monkey, Portland, USA). Ethical approval was provided by Queen Mary University of London (QMUL) ethics committee (Ref: QMERC2014/76).

Participants: An opportunistic sample of UK resident post-natal mothers with a self-declared diagnosis of GDM was recruited via on-line forums such as Diabetes UK forum, Netmums and Mumsnet, and social media. Eligibility was confirmed through explicit questions within the survey. The healthcare professionals were identified via email and social media at Barts Health NHS Trust. “Snowballing” was encouraged, whereby respondents could forward the online survey link to their contacts, to maximise the diversity of response¹⁵. Only UK healthcare professionals who supported postnatal women as part of their normal clinical

practice, confirmed by explicit questions. Respondents were excluded if they did not meet these criteria.

Survey: The 28-item questionnaire for mothers was designed for this project to develop a understanding of what type of intervention mothers who had GDM would find helpful and engaging. The design was informed by conducting post-natal discussion groups with mothers who were diagnosed with GDM in their recent pregnancy. Questions covered recruitment timing, intervention content, format and delivery as well as health behaviour and attitudes to diet, weight and exercise. Basic demographic data were also collected. The healthcare professionals' survey comprised of 17 questions focussing on perceived barriers and facilitators to discuss weight-related issues in the postnatal period in women with GDM, their current practice to support these women, views about what would make an intervention successful, appropriate timing of intervention in the postpartum period, and their interpretation of their patient's motivation to engage in lifestyle changes to minimise risk of type 2 diabetes.

The questionnaires included both open format questions, multiple choice questions and questions with a Likert scale response¹⁶. Both questionnaires were piloted within a small sample to check the clarity and performance of the on-line questionnaire.

Analysis:

Participants' response to individual questions were reported as a median score and analysed using Mann-Whitney and Kruskal Wallis non-parametric tests^{16,17}. Certain sets of questions, such as those concerning general attitudes to exercise, weight and diet on the mothers' survey were analysed collectively as Likert Scale data, to give a composite health behaviour score which was analysed by ANOVA^{16,17}. Scoring of negative questions was reversed, so a high score was an indicator of positive attitude towards postnatal health behaviour. We grouped ethnicity as Caucasian, and Black and minority ethnic group (BME), Body Mass Index as

normal weight 18-24.99; overweight 25-29.99; obese ≥ 30 , and educational status as highest attainment GCSE, A level or tertiary education. All analyses were undertaken using SPSS 23.0 (SPSS Inc., Chicago, USA). Completed surveys were checked for missing data and responses in which a large number of questions were skipped, were excluded from the analysis.

Results

Overall 129 participants completed the surveys, including 83 mothers and 46 healthcare professionals.

Participant characteristics

The characteristics of mothers who participated are provided in Table 1. Participants were predominantly Caucasian (67%) and well educated (64% had tertiary education). Fifty eight percent were either overweight or obese and 17% were normal weight. 25% of participants chose not to provide their weight and height. The mean BMI was 28.38 (SD 5.04). The sample of healthcare practitioners comprised of 24 obstetricians, 8 diabetologists, 8 midwives, 5 General Practitioners, and 1 dietitian. They were involved in the care of mothers from a predominantly South Asian population.

Design of a lifestyle intervention

When is the best time to engage women?

The responses from mothers varied on optimal timing to receive information on lifestyle and weight management, with 26% (21/81) preferring six-weeks, 23% (19/81) in the first three months, 31% (25/81) between 3 and 6 months and 20% (16/81) after 6 months. Fifty-two

percent (24/46) of professionals felt that the 6-week check was the ideal time to engage women and 48% (22/46) felt it should be flexible according to when a mother was ready.

How should the intervention be delivered?

The preferred choice of delivery of the intervention or service was group sessions (30%, 25/83) followed by mobile/internet support (24%, 20/83) and 1:1 support sessions (24%, 20/83). Home visits and a buddy scheme were less popular modes of intervention delivery, (13% and 9% respectively (11/83 and 7/83)). Mothers were more likely engage in sessions held in the community (39%, 32/83) than in a medical setting (15%, 12/83). Group sessions were also recommended by 45% (19/42) of professionals; 1:1 sessions with a specialist were preferred by 25% (10/42), a buddy scheme by 12% (5/42) mobile/internet support by 9% (4/42) and home visits by 9% (4/42).

What advice should the intervention programme include?

Mothers felt key elements of a programme to promote a healthy lifestyle and weight for postpartum women with GDM included an exercise programme (95%, 79/83), recipe ideas (85%, 71/83) and dietary advice (76%, 63/83). Sixty-one per cent (51/83) supported a weekly weigh-in session. Cooking skills sessions were the least popular element selected, in this sample of women (39%, 32/83).

Health care professionals considered dietary advice and an exercise programme to be the most important elements to include (92% and 91% respectively). Sixty-five percent (27/42) supported the inclusion of a weekly weigh in, 62% (26/42) recipe ideas and 57% (24/42) cooking skills sessions (Fig 4).

Mothers were less in favour of including cooking sessions ($P=0.043$) and general dietary advice ($P=0.012$) than were health professionals and preferred recipe ideas ($P<0.001$). (Fig. 4.).

Health behaviour and barriers towards optimal health behaviours amongst post-natal GDM mothers

When asked *Which of these factors, if any, is most important in reducing the risk of developing type 2 diabetes after GDM?* 13% said eating healthily, 7% maintaining a healthy weight, 2% exercising regularly and 78% said all of these factors were important.

Overall, the respondents followed the dietary advice they were given in pregnancy (81% (67/83) agreed, Median 4, IQR 1) and in the postnatal period (65% (54/83) agreed, Md 4, 1) and felt they knew what they should be eating (81% (67/83) disagreed with the statement *I don't know what should be included in a healthy diet*, Md score 1, IQR 1). Mothers did not feel that time to prepare meals or the cost of healthy foods, were barriers to eating healthily (62% (51/83) disagreed that *cooking healthily takes too long*, Md score 2, IQR 1; 57% (48/83) disagreed with the statement *healthy food is too expensive*, Md score 2, IQR 2). Forty three percent (36/83) did not think it was too hard to think about eating healthily when you are looking after a new baby, (Md score 3, IQR 2) but 38% (31/83) felt it was. (Fig. 1).

Barriers to exercise included tiredness (*“Women are too tired to exercise when they have a young baby”* - 65% (54/83) agreed, Md score 4, IQR 1) and the need for childcare (*“it is hard to exercise in the postnatal period because of the need for childcare”* (64% agreed (53/83), Md score 4, IQR 2). Seventy per cent of respondents (57/81) cited walking as a normal part of their daily routine as their main form of physical activity, 9% (7/81) walked specifically for exercise, 6% (5/81) did not exercise at all, and the remainder took part in a mixture of activities from organised sport, exercising to DVDs at home and using local sports facilities

such as gyms and exercise classes. The most commonly chosen factors that would help participation in exercise were free classes (36%, 29/81) and free childcare (33%, 27/81). Younger participants (under 35) had a more positive attitude to physical activity than older mothers ($P=0.04$) (Fig. 2). Losing weight was a concern for 69% (57/83) of the women (Md 4, IQR 1), 69% (57/83) felt it was important to aim to return to their pre-pregnancy weight (Md 4, IQR 2). Women who were overweight or obese were more likely to be concerned about losing weight than those of normal weight ($P=0.03$). Forty-one percent (34/83) of mothers agreed it was hard to focus on weight with a new baby (Md 3, IQR 2). Seventy-six per cent (63/83) would like more information about managing their weight in the postnatal period (Md 4, IQR 0). (Fig.3). Women were most likely to seek help to lose postnatal weight from friends and family (46%, 38/83) or a support group in the community (39%, 32/83). Thirty-six per cent said they would seek help from their GP, 23% (30/83). Forty-one per cent (34/83) said they would be more comfortable talking to a female health care practitioner, rather than a male, about their weight (Md 4, IQR 2) and this was not associated with ethnicity ($P=0.287$). (Fig.3).

The composite score from all the attitudinal health behaviour questions indicated a positive attitude towards postnatal health behaviour (mean 3.50, SD 0.39). This was not associated with ethnicity ($P=0.117$), BMI ($P=0.272$), Mothers' age ($P=0.317$) or mother's education ($P=1.121$). Mothers placed equal importance on the importance of exercise, weight management and physical activity ($P=0.268$).

Health professionals

Fifty-six percent of health professionals (26/46) did not think that women were aware of the importance of controlling their weight in reducing future risk of type 2 diabetes (Md score 2, IQR 1). Thirty-nine percent of health professionals (18/46) found it difficult to raise the issue

of weight with postnatal women who were diagnosed with GDM (Md score 3, IQR 2), compared to 59% (27/46) when advising pregnant women with risk factors for GDM, (*“I know I should talk about weight but sometimes it’s a difficult subject to raise”*) (Md score 4, IQR 2). Eighty-three per cent (38/46) felt it was *“vital”* they raised the issue of weight in postnatal women who had been diagnosed with GDM (Md score 4, IQR 1). Healthcare practitioners agreed they would like to be involved in supporting women with postnatal weight management (69% (29/42) agreed, Md 4, IQR 1) and 41% (17/42) currently felt they were equipped to do so (Md 3, IQR 2). However, 83% (35/42) felt they did not have enough time to help postnatal women with weight-loss, in their day-to-day roles (md 4, IQR 0). Seventy-two percent of health professionals felt that exercising for health was not a part of their patients’ regular behaviour. Perceived engagement in exercise was not associated with patient ethnicity in this sample ($P = 0.082$) although 87% (20/23) of healthcare practitioners dealing with BME patients felt exercise was not part of their patients’ culture compared to only 57% (4/7) of those with a primarily Caucasian patient base.

Discussion

This survey was conducted to inform the design of a lifestyle intervention for postpartum women who had gestational diabetes in pregnancy, as per MRC guidance for complex interventions¹⁸. It provides novel data from a sample of UK postnatal mothers who had GDM and health professionals, about what they recommend for a postnatal lifestyle intervention to reduce the risk of the subsequent development of type 2 diabetes risk. Women in this sample were supportive of a group-based lifestyle programme offered in the community that included recipe ideas, an exercise programme and weekly weigh-in sessions. However, mothers also indicated significant barriers to engagement are present in the postnatal period.

This sample of women indicated that they were aware that eating healthily, keeping physically active and managing their weight are important in reducing their subsequent diabetes risk. However, a gap between knowledge and behaviour has been reported in the GDM postpartum population^{19, 20}, as well as a poor perception of personal risk which may limit personal behaviour change^{21, 22}. More than half of the health practitioners in caring for GDM women in this survey, do not feel that women are aware of the importance of controlling their postpartum weight to reduce the risk of developing type 2 diabetes. Reducing postpartum weight retention is a key focus of postpartum lifestyle intervention; in women diagnosed with GDM, postpartum weight gain of 4.5kg, doubled the risk of subsequently developing diabetes within the 7.5 year follow-up^{23, 24}. A modest post-natal weight-loss (anything more than 2kg) is associated with improved glucose metabolism at 12 months post-partum in women with GDM²⁵. Studies have suggested that health practitioners' confidence in raising the issue of weight, may be a barrier to engaging obese pregnant women in managing their weight^{26, 27}. Although healthcare practitioners in this survey also felt it was sometimes difficult to raise the subject, they considered this to be a “*vital*” part of their job.

Both healthcare practitioners and mothers agreed that exercise, dietary support were important elements in a postnatal lifestyle intervention for GDM mothers. A recent Cochrane review of lifestyle interventions after childbirth, supports that the most effective interventions for weight-loss include either diet or diet and exercise together²⁸. Women in this sample wanted inspiration in the form of recipe ideas, rather than specific dietary advice or cooking skills support. The difference in the value placed on improving knowledge and cooking skills between the healthcare practitioners and mothers in this survey, may well be due to the demographic differences between the health practitioner patient base and mothers in this sample. Women in this study recognised the importance of physical activity although analysis

of a postpartum GDM cohort in Sweden suggested that in practice the postpartum physical activity level of GDM mothers may be lower than in mothers who had a normo-glycaemic pregnancy²⁰. Results from the postpartum KAN-DO intervention trial, where women were overweight or obese when they entered pregnancy, showed very small increases in reported physical activity as a result of intervention, suggesting that tailored interventions are needed to help overweight postpartum women integrate physical activity and reduce sedentary behaviour²⁹.

While this study provides valuable insight into the acceptability of intervention and possible content and setting, we were limited by the small sample and self-selection of participants, affecting the representativeness of the sample. The mothers were primarily Caucasian and well-educated, which may limit how applicable the results are to low-income ethnic minority women who are particularly at risk. While health professionals who support women through pregnancy and their gestational diabetes management are well represented in this study, we had limited response from GPs and health visitors who may have provided further insight into the challenges faced by postpartum mothers. We were not able to calculate participation rate nor compare participants to non-participants as questionnaires were posted on online forums rather than targeted at a predetermined group. In addition, we are not able to comment on whether any differences in responses between Internet users and non-Internet users might exist.

Cultural acceptability of exercise varies it is recognised that it is less acceptable for women in South Asian communities in particular³⁰. The health professionals' results indicate an awareness of cultural constraints towards exercise in ethnic minority patients. As BME women of reproductive age, are particularly at risk of obesity³¹ and GDM³², they are a key target for lifestyle interventions. Care therefore needs to be given to these cultural sensitivities when designing lifestyle programmes. The exercise element of lifestyle

intervention is most likely to be successful when objectively designed goals are used, for example 10,000 steps with a pedometer or use of heart-rate monitors³³. Results from this survey suggest a focus on the inclusion of walking targets would be acceptable and achievable way to promoting an increase in exercise.

Postpartum women have proved hard to recruit and to engage in postnatal lifestyle interventions³⁴⁻³⁶. Our survey supports current literature that suggests looking after a new baby makes it difficult for mothers to focus on their health; lack of time to cook or exercise, lack of family support, low self-esteem and mood issues have all been shown to be negative influences^{9, 37}. Postpartum fatigue may also be present for a number of months after delivery³⁸. Timing of recruitment of postpartum mothers is likely to be an important consideration to maximise recruitment and retention of mothers in a lifestyle intervention³⁵. Meta-analysis of behavioural change techniques used in lifestyle interventions, indicate that self-monitoring is a key strategy in intervention success, particularly if combined Control Theory such as specific goal setting or feedback on performance³⁹. Encouraging women to regularly self-monitor and record their weight is a simple promising strategy⁴⁰ and has been validated for GDM mothers⁴¹.

Pragmatic issues such as optimising convenience, timings of sessions and tailoring of the programme to personal needs have all been shown to be valuable strategies to improve engagement³⁵. The acceptability of group sessions should be further explored. Home-based elements of support within an intervention have the capacity to be cost-effective and wide reaching compared to intensive face-to-face counselling⁴². The use of technology-based intervention needs further evaluation of the accessibility and appeal in the BME populations.

Summary and implications

Post-GDM mothers in this sample reported good knowledge of the health behaviour that would reduce their subsequent diabetes risk. They recommended that a lifestyle intervention should be group-based and offered in the community, although reported barriers of time constraints, child care issues and difficulty prioritising health, suggests that flexible options for engagement should be included– such as internet support and flexible sessions with inclusion of children or a childcare facility. While some were ready to engage relatively soon after giving birth, others would prefer lifestyle advice to be available when they were ready, suggesting a flexible window for recruitment may be beneficial.

Funding sources: No funding was received for this study.

Conflicts of interest: None

Acknowledgements: The authors would like to thank Diabetes UK, Netmums and Mumsnet for hosting a link to the survey.

References

1. Bellamy L, Casas JP, Hingorani AD, Williams D. Type 2 diabetes mellitus after gestational diabetes: a systematic review and meta-analysis. *Lancet*. May 23 2009;373(9677):1773-1779.
2. Kim C, Newton KM, Knopp RH. Gestational diabetes and the incidence of type 2 diabetes: a systematic review. *Diabetes Care*. Oct 2002;25(10):1862-1868.
3. Beharier O, Sergienko R, Kessous R et al. "Gestational diabetes mellitus is a significant risk factor for long-term ophthalmic morbidity." *Archives of Gynecology and Obstetrics* Jun 2017;295(6):1477-1482.
4. Fuchs O, Sheiner E, Meirovitz M et al. "The association between a history of gestational diabetes mellitus and future risk for female malignancies." *Archives of gynecology and obstetrics* Mar 2017 295(3): 731-736.5. Harreiter J, Dovjak G, Kautzky-Willer A. Gestational diabetes mellitus and cardiovascular risk after pregnancy. *Womens Health (Lond Engl)*. Jan 2014;10(1):91-108.
6. Aroda VR, Christophi CA, Edelstein SL, et al. The effect of lifestyle intervention and metformin on preventing or delaying diabetes among women with and without gestational diabetes: The diabetes prevention program outcomes study 10-year follow-up. *Journal of Clinical Endocrinology and Metabolism*. 2015;100(4):1646-1653.

7. Morton S, Kirkwood S, Thangaratinam S. Interventions to modify the progression to type 2 diabetes mellitus in women with gestational diabetes: a systematic review of literature. *Curr Opin Obstet Gynecol*. Dec 2014;26(6):476-486.
8. Guo J, Chen JL, Whittemore R, Whitaker E. Postpartum Lifestyle Interventions to Prevent Type 2 Diabetes Among Women with History of Gestational Diabetes: A Systematic Review of Randomized Clinical Trials. *J Womens Health (Larchmt)*. Jan 2016;25(1):38-49.
9. Montgomery KS, Bushee TD, Phillips JD, et al. Women's challenges with postpartum weight loss. *Maternal and Child Health Journal*. Nov 2011;15(8):1176-1184.
10. Onyango AW, Nommsen-Rivers L, Siyam A, et al. Post-partum weight change patterns in the WHO Multicentre Growth Reference Study. *Matern Child Nutr*. Jul 2011;7(3):228-240.
11. Walker LO, Sterling BS, Latimer L, Kim SH, Garcia AA, Fowles ER. Ethnic-specific weight-loss interventions for low-income postpartum women: findings and lessons. *West J Nurs Res*. Aug 2012;34(5):654-676.
12. O'Dea A, Tierney M, McGuire BE, et al. Can the onset of type 2 diabetes be delayed by a group-based lifestyle intervention in women with prediabetes following gestational diabetes mellitus (GDM)? Findings from a randomized control mixed methods trial. *Journal of Diabetes Research*. 2015;2015.
13. Morrison MK, Lowe JM, Collins CE. Perceived risk of Type 2 diabetes in Australian women with a recent history of gestational diabetes mellitus. *Diabet Med*. Aug 2010;27(8):882-886.
14. Ferrara A, Ehrlich SF. Strategies for diabetes prevention before and after pregnancy in women with GDM. *Current Diabetes Reviews*. 2011;7(2):75-83.
15. Sixsmith J, Boneham M, Goldring JE. Accessing the community: gaining insider perspectives from the outside. *Qualitative Health Research*. Apr 2003;13(4):578-589.
16. Johns R. Likert Items and Scales: Survey Resource Network; 2010.
17. Boone H, Boone D. Analysing likert data. *Journal of Extension*. Vol 50; 2012.
18. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Int J Nurs Stud*. May 2013;50(5):587-592.
19. Jones EJ, Roche CC, Appel SJ. A review of the health beliefs and lifestyle behaviors of women with previous gestational diabetes. *JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*. 2009;38(5):516-526.
20. Persson M, Winkvist A, Mogren I. Lifestyle and health status in a sample of Swedish women four years after pregnancy: A comparison of women with a history of normal pregnancy and women with a history of gestational diabetes mellitus. *BMC Pregnancy and Childbirth*. 2015;15(1).
21. Kieffer EC, Carman WJ, Gillespie BW, Nolan GH, Worley SE, Guzman JR. Obesity and gestational diabetes among African-American women and Latinas in Detroit: implications for disparities in women's health. *J Am Med Womens Assoc*. Fall 2001;56(4):181-187, 196.
22. Mukerji G, Kainth S, Pendrith C, et al. Predictors of low diabetes risk perception in a multi-ethnic cohort of women with gestational diabetes mellitus. *Diabet Med*. Oct 2016;33(10):1437-1444.

23. Walker LO. Managing excessive weight gain during pregnancy and the postpartum period. *J Obstet Gynecol Neonatal Nurs.* Sep-Oct 2007;36(5):490-500.
24. Peters RK, Kjos SL, Xiang A, Buchanan TA. Long-term diabetogenic effect of single pregnancy in women with previous gestational diabetes mellitus. *Lancet.* Jan 27 1996;347(8996):227-230.
25. Ehrlich SF, Hedderson MM, Quesenberry CP, Jr., et al. Post-partum weight loss and glucose metabolism in women with gestational diabetes: the DEBI Study. *Diabetic Medicine.* Jul 2014;31(7):862-867.
26. Knight BA, Wyatt K. Barriers encountered when recruiting obese pregnant women to a dietary intervention. *Nursing Times.* Aug 17-23 2010;106(32):20-22.
27. Heslehurst N, Newham J, Maniatopoulos G, Fleetwood C, Robalino S, Rankin J. Implementation of pregnancy weight management and obesity guidelines: a meta-synthesis of healthcare professionals' barriers and facilitators using the Theoretical Domains Framework. *Obes Rev.* Jun 2014;15(6):462-486.
28. Amorim AR, Linne YM, Lourenco PM. Diet or exercise, or both, for weight reduction in women after childbirth. *Cochrane Database Syst Rev.* 2007(3):CD005627.
29. Evenson KR, Brouwer RJN, Østbye T. Changes in Physical Activity Among Postpartum Overweight and Obese Women: Results from the KAN-DO Study. *Women and Health.* 2013;53(3):317-334.
30. Vyas A, Chaudhary N, Ramiah K, Douglass K. Risk perceptions, nutrition and physical activity among South Asian women in the US: does history of gestational diabetes mellitus (GDM) matter? *Health.* 2012;4(12):1263-1270.
31. Gatineau M, Mathrani S. Ethnicity and obesity in the UK. *Perspectives in Public Health.* Jul 2011;131(4):159-160.
32. Poston L, Bell R, Croker H, et al. Effect of a behavioural intervention in obese pregnant women (the UPBEAT study): a multicentre, randomised controlled trial. *Lancet Diabetes Endocrinol.* Oct 2015;3(10):767-777.
33. Nascimento SL, Pudwell J, Surita FG, Adamo KB, Smith GN. The effect of physical exercise strategies on weight loss in postpartum women: a systematic review and meta-analysis. *International Journal of Obesity.* 2014;38(5):626-635.
34. Ferrara A, Hedderson MM, Albright CL, et al. A pregnancy and postpartum lifestyle intervention in women with gestational diabetes mellitus reduces diabetes risk factors: a feasibility randomized control trial. *Diabetes Care.* Jul 2011;34(7):1519-1525.
35. Atkinson L, Olander E, French D. Why don't many obese pregnant and post-natal women engage with a weight management service? *Journal of Reproductive and Infant Psychology.* 2013;31(3):245-256.
36. Carter-Edwards L, Ostbye T, Bastian LA, Yarnall KS, Krause KM, Simmons TJ. Barriers to adopting a healthy lifestyle: insight from postpartum women. *BMC Research Notes.* 2009;2:161.
37. Pedersen P, Baker JL, Henriksen TB, et al. Influence of psychosocial factors on postpartum weight retention. *Obesity (Silver Spring).* Mar 2011;19(3):639-646.
38. Troy NW. Is the significance of postpartum fatigue being overlooked in the lives of women? *MCN Am J Matern Child Nurs.* Jul-Aug 2003;28(4):252-257.

39. Michie S, Abraham C, Whittington C, McAteer J, Gupta S. Effective techniques in healthy eating and physical activity interventions: a meta-regression. *Health Psychology*. Nov 2009;28(6):690-701.
40. van der Pligt P, Willcox J, Hesketh KD, et al. Systematic review of lifestyle interventions to limit postpartum weight retention: implications for future opportunities to prevent maternal overweight and obesity following childbirth. *Obesity Reviews*. Oct 2013;14(10):792-805.
41. Paez KA, Griffey SJ, Thompson J, Gillman MW. Validation of self-reported weights and heights in the avoiding diabetes after pregnancy trial (ADAPT). *BMC Medical Research Methodology*. 2014;14(1).
42. van der Pligt P, Willcox J, Hesketh KD, et al. Systematic review of lifestyle interventions to limit postpartum weight retention: implications for future opportunities to prevent maternal overweight and obesity following childbirth. *Obes Rev*. Oct 2013;14(10):792-805.

Figures

Fig 1




Fig 2




Fig 3




Fig 4




Table 1: Sample characteristics of the mothers who participated (n=83)

	n (%)	Mean \pm sd
Age		34.8 \pm 5.0
BMI		28.38 \pm 5.04
BMI Category		
Normal weight	14 (17)	
Overweight/obese	48 (58)	
Unknown	21 (25)	
Educational Attainment		
GCSE	4 (5)	
A Level or equivalent	26 (31)	
Tertiary education	53 (64)	
Ethnicity		
Caucasian	56 (67)	
BME	27 (33)	
Age of youngest child		
Under 1	33 (40)	
1-2 yrs	20 (24)	
3+ yrs	30 (36)	